

Interns Overestimate the Effectiveness of Their Hand-off Communication

AUTHORS: Vivian Y. Chang, MD,^a Vineet M. Arora, MD, MA,^b Shiri Lev-Ari, MA,^c Michael D'Arcy, BA,^d and Boaz Keysar, PhD^c

^aDivision of Pediatric Hematology/Oncology, University of California, Los Angeles, Los Angeles, California; Departments of ^bMedicine and ^cPsychology, University of Chicago, Chicago, Illinois; and ^dDepartment of Medical Anthropology, University of California, Berkeley, Berkeley, California

KEY WORDS

work hours, hand-off communication, patient safety

www.pediatrics.org/cgi/doi/10.1542/peds.2009-0351

doi:10.1542/peds.2009-0351

Accepted for publication Sep 28, 2009

Address correspondence to Vineet M. Arora, MD, MA, 5841 S Maryland Ave, MC 2007, AMB B217, Chicago, IL 60637. E-mail: varora@medicine.bsd.uchicago.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2010 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: *Dr Chang was a paid resident of the University of Chicago Comer Children's Hospital at the time of this study; Dr Arora is a paid employee of the University of Chicago Medical Center, which trains interns and residents; Ms Lev-Ari, Mr D'Arcy, and Dr Keysar have no financial relationships relevant to this article to disclose.*



WHAT'S KNOWN ON THIS SUBJECT: Transitions of patient care from 1 physician to another, otherwise known as hand-offs, are riddled with omitted or inaccurate information, leading to near-misses or adverse events. It is unclear why physician communication is so problematic.



WHAT THIS STUDY ADDS: This study found that pediatric residents overestimated the effectiveness of their communication during hand-offs. Theories from the psychology of communication propose that such overestimation is a systematic cause of miscommunication.

abstract

OBJECTIVE: Theories from the psychology of communication may be applicable in understanding why hand-off communication is inherently problematic. The purpose of this study was to assess whether postcall pediatric interns can correctly estimate the patient care information and rationale received by on-call interns during hand-off communication.

METHODS: Pediatric interns at the University of Chicago were interviewed about the hand-off. Postcall interns were asked to predict what on-call interns would report as the important pieces of information communicated during the hand-off about each patient, with accompanying rationale. Postcall interns also guessed on-call interns' rating of how well the hand-offs went. Then, on-call interns were asked to list the most important pieces of information for each patient that postcall interns communicated during the hand-off, with accompanying rationale. On-call interns also rated how well the hand-offs went. Interns had access to written hand-offs during the interviews.

RESULTS: We conducted 52 interviews, which constituted 59% of eligible interviews. Seventy-two patients were discussed. The most important piece of information about a patient was not successfully communicated 60% of the time, despite the postcall intern's believing that it was communicated. Postcall and on-call interns did not agree on the rationales provided for 60% of items. In addition, an item was more likely to be effectively communicated when it was a to-do item (65%) or an item related to anticipatory guidance (69%) compared with a knowledge item (38%). Despite the lack of agreement on content and rationale of information communicated during hand-offs, peer ratings of hand-off quality were high.

CONCLUSIONS: Pediatric interns overestimated the effectiveness of their hand-off communication. Theories from communication psychology suggest that miscommunication is caused by egocentric thought processes and a tendency for the speaker to overestimate the receiver's understanding. This study demonstrates that systematic causes of miscommunication may play a role in hand-off quality. *Pediatrics* 2010; 125:491–496

In 2003, the Accreditation Council for Graduate Medical Education restricted residency duty hours in response to increasingly complicated patients, data on the effects of sleep deprivation on residents, and growing media attention on the affects of long duty hours on patient safety and resident well-being.¹ Since then, many voiced concerns about the increased frequency of hand-offs, whereby patient care responsibility is transferred from 1 resident who is leaving the hospital to another resident who will be staying and covering other residents' patients.^{2,3} The frequency of hand-offs did in fact increase after the duty hour restrictions, with a member of the primary health care team being present in the hospital for less than half of a patient's hospitalization.⁴ This emphasizes the importance of quality hand-offs.

A recent *Pediatrics* article⁵ found no significant change in the total hours of work or sleep before and after the duty hour restrictions. It is interesting that the authors found a significant increase in minor errors, which may result from an increase in hand-off frequency without a corresponding increase in hand-off education and improvement. In addition, in December 2008, the Institute of Medicine released recommendations to reduce work hours further and to train residents on transitions of patient care.⁶ Implementing these recommendations would further increase the frequency of hand-offs. Thus, undertaking formal hand-off education and improvement activities is becoming increasingly critical to ensuring safe patient care. To develop such programs, it is essential to understand the systemic reasons for hand-off communication failure.

The hand-off process, also known as "sign-out," can be a written or verbal transfer of patient care information. Each time a hand-off occurs, the possi-

bility for miscommunication arises. Hand-offs are often riddled with omitted or inaccurate information that could be critical to patient care, such as code status or allergies, resulting in uncertainty in the covering residents' decisions for patients.^{7,8} The contribution of communication failures to adverse events has been estimated to be between 15% and 67%.⁹⁻¹¹ To date, little is known about the hand-off process in general and in the field of pediatrics in particular. Although several studies focused on near-misses and adverse events,^{7,12,13} they lacked the theoretical foundation to explain why physician-to-physician communication is poor.

Theories from the psychology of communication may be applicable in understanding why hand-off communication is inherently problematic. Studies show that speakers systematically overestimate how well their messages are understood by listeners¹⁴ and that people in general believe that their thoughts are transparent to others.¹⁵⁻¹⁷ In addition, the more knowledge that people share, the worse they communicate new material because they overestimate the knowledge of the other.¹⁸ These psychological processes could systematically affect the effectiveness of communication during hand-offs. The aim of this study was to assess whether postcall pediatric interns who provide hand-offs can correctly estimate the information received by on-call interns at a hand-off communication.

METHODS

Participant Population

All interns, subinterns, and visiting interns who were rotating on the general pediatrics team at the University of Chicago Comer Children's Hospital were eligible for the study. During June 2007, the study protocol was explained by pediatric resident and investigator (Dr Chang), and written con-

sents were obtained. Participation was voluntary, and the institutional review board at the University of Chicago approved this study.

Study Setting


At the University of Chicago, there is 1 general pediatrics team that is composed of 1 attending, 2 senior residents (postgraduate year 3) who provide day coverage from 7 AM until 7 PM, 2 senior residents (postgraduate year 2 or 3) who provide night coverage from 7 PM until 7 AM, and 4 interns (postgraduate year 1). Interns rotate through general pediatrics for 1.5 to 2.0 months. One intern is on call each night and works 30-hour shifts every 4 days. The on-call intern admits general pediatric patients from 7 AM to 7 AM the next day. The verbal hand-off occurs at 11:30 AM daily in a dedicated conference room with the postcall intern communicating primarily to the on-call intern but with the other interns and senior residents also present. Senior residents are present in the room mainly to play a supervisory role and could interject if necessary. The attending is not present. The computerized written hand-off is a Microsoft Word document (Redmond, WA) updated by the postcall intern and given to the on-call intern.

Data Collection

Participant Recruitment

Interns were interviewed on the last 4 weekdays of their general pediatrics rotation. This allowed interns to establish their own hand-off practice patterns, minimizing the potential for the interview to act as an intervention, or the Hawthorne Effect.¹⁹

On-call and postcall interns were approached after the hand-off and asked to participate in an interview about the hand-off communication. The postcall intern was interviewed immediately after the handoff, allowing that intern to

**FIGURE 1**

Timing of interviews. Hand-offs occurred daily at 11:30 AM. The postcall intern was interviewed immediately after the hand-off, and the on-call intern was interviewed at 1:00 PM after and educational noon conference.

leave the hospital within duty hour restrictions. The on-call intern was interviewed after a noon conference, at 1:00 PM (Fig 1). The interns had access to the written hand-off during the interview, which took place in a private room and was audiotaped. All interviews were conducted by trained research assistants (Ms Lev-Ari and Mr D'Arcy).

Postcall Interview Script

The interviewer asked the postcall interns to identify patients by room number to protect health information. They were told that the on-call intern will be asked to describe the most important pieces of information for each patient

that was communicated during the hand-off and then asked to rank the information in order of importance (Table 1). The interviewer also explained that the on-call intern will be asked to report any rationale that he or she received for each item. The interviewer then asked the postcall interns to detail how they expect the on-call intern to answer these questions. This provided an estimate of what the postcall intern believed that the on-call intern received from the hand-off session.

The postcall intern was then asked to guess the on-call intern's rating of how well the hand-off prepared him or her to take care of the patients, on a scale

from 1 to 10, with 1 being "did not prepare me at all" and 10 being "prepared me very well." Finally, the postcall intern guessed the on-call intern's rating of how well the hand-off went in general, on a scale from 1 to 10, 1 being "badly" and 10 being "very well."

On-call Interview Script

The interviewer referred to patients by the room numbers provided by the postcall intern. The interviewer asked the on-call intern to list the most important pieces of information, along with any rationale, about each patient that the postcall intern discussed during the hand-off and then to rank this information in order of importance. These questions were repeated for each patient. This provided an estimate of what the on-call intern actually received from the handoff session. Then, the on-call interns rated how well the hand-off prepared them to take care of the patients overnight and how well the hand-off went in general, both on a scale from 1 to 10.

Data Analysis

Audiotaped interviews were transcribed by Voss Transcription, Inc (Chicago, IL), and reviewed by 2 physician investigators (Drs Chang and Arora) for accuracy. Occasionally, when transcripts had missing information, the original audiotapes were consulted by

TABLE 1 Interview Script

Postcall Intern	On-call Intern
We will ask the on-call physician to report what information he or she received about this patient, as well as the rationale you provided for each piece of information. We will ask him or her to list the pieces of information, starting with the most important information. We would like to ask you to guess how he or she answered, starting with the most important piece of information.	Could you please tell me what information you received about this patient? For each piece of information, please include the rationale that the primary physician provided when possible. Please list the pieces of information in order of importance, starting with the most important information.
We will ask the on-call physician to rate how well the sign-out went, with 1 being "badly" and 10 being "very well." Could you guess his or her rating, and can you please explain your answer?	Could you rate how well you feel the sign-out went, with 1 being "badly" and 10 being "very well"? Could you please explain your rating?
We will ask the on-call physician to rate how well the sign-out prepared him or her to take care of these patients, with 1 being "did not prepare me at all" and 10 being "it prepared me very well." Could you guess his or her rating, and can you briefly explain your answer?	Could you rate how well you feel the sign-out prepared you to take care of these patients, with 1 being "did not prepare me at all" and 10 being "it prepared me very well"? Could you briefly explain your rating?

Postcall interns were asked to guess the on-call interns' responses to interview questions. On-call interns were asked to list the important items communicated during the hand-off about each patient and accompanying rationale. They rated how well the hand-off went and how well-prepared they felt to take care of patients overnight.

physician investigators (Drs Chang and Arora) and blanks were filled in when possible. For example, “sickle-dex” was inaudible to transcribers, but given the investigators’ knowledge of the hospital’s qualitative screen for sickle hemoglobin, the missing name for the laboratory test was filled in.

Transcripts were stripped of identifiers. Postcall and on-call transcripts for the same hand-off were paired. For each patient, data were extracted into a Microsoft Excel spreadsheet, and important items that the postcall and on-call interns reported were juxtaposed. The items were then coded for agreement between the interns, with 1 being “agreement” and 0 being “no agreement.” To qualify as agreement, the item had to match in meaning, not necessarily in terminology. We then compared the rationale that the postcall intern expected the on-call intern to report with the rationale that the on-call intern actually reported. Last, we coded items with respect to importance: (1) the most important item about each patient and (2) the top 3 important items regardless of priority rank. The coding was initially conducted by 1 investigator (Dr Chang) and then reviewed by a second investigator (Dr Arora).

We used summary statistics to tabulate the mean agreement on the most important items about each patient, how well the hand-off prepared the on-call intern to take care of patients, and how well the overall hand-off went. We conducted subgroup analyses by using χ^2 tests for type of intern and type of item being communicated. All statistical tests were performed by using Stata 10.0 (College Station, TX), with statistical significance defined as $P < .05$.

RESULTS

All 18 categorical pediatric interns and 5 combined medicine-pediatric interns (100%) agreed to participate in the

study from July 2007 to May 2008. Ten of 28 visiting interns (family medicine interns rotating from other hospitals) or subinterns (fourth-year medical students from the study institution) also agreed to participate. We conducted 52 interviews, which constituted 59% of eligible interviewees. Of the 52 interviews, there were 19 dyad interviews, consisting of both the postcall and the on-call interns of the same hand-off. The majority of interviews that were included in the analysis were from categorical pediatric interns (63%). Sixteen percent of the analyzed interviews were from medicine-pediatric interns, 18% were from rotating interns, and only 1 was from a subintern (<2%). Fourteen interviews were discarded because only 1 of the 2 interns was interviewed and therefore could not be paired. Seventy-two patients were discussed during the interviews that were analyzed.

Postcall interns overestimated the effectiveness of their communication. For example, 1 postcall intern expected the on-call intern to have understood the following about a patient:

1. “Follow-up on surgery’s recommendations.”
2. “Postop, restart patient on feeds and if that improves, stop [intravenous] fluids.”
3. “Patient will stay on [intravenous] antibiotics today and will go by mouth tomorrow.”

In contrast, the on-call intern actually understood:

1. “Coming back from surgery, so restart feeds.”
2. “I might get a page from [affiliated hospital] and I’ll just defer to primary physician.”

This on-call intern mentioned only 1 of the 3 items that the postcall intern expected. This discrepancy was very common. On average, postcall interns expected on-call interns to mention 2.6

important items per patient, whereas on-call interns actually mentioned only 1.6 items on average ($P < .01$). For 69% of the patients, the on-call intern failed to note at least 1 of the important items that the postcall intern expected him or her to note.

We also looked at how interns ranked the items about each patient, in order of importance, as a measure of how well the gravity of each item was communicated. The postcall interns overestimated their ability to convey the information about the importance of each item. Overall, the item that postcall interns expected on-call interns to perceive as the most important was not perceived as such by the on-call interns for 60% of the patients. In fact, the most important item about a patient was not mentioned at all by the on-call intern for 40% of the patients.

We conducted subset analyses comparing categorical pediatric interns, combined medicine-pediatric interns, and rotating family medicine interns. There was no difference between categorical pediatric interns and combined medicine-pediatric interns in the percentage of items that were successfully communicated during the hand-off; however, when the postcall intern was a rotating family medicine intern, there was a significantly lower likelihood that the most important item about a patient was communicated (odds ratio: 0.16 [95% confidence interval: 0.04–0.75]; $P = .02$). In addition, the percentage of overall agreement was significantly lower compared with pediatric interns (95% confidence interval: 9.1%–49.0%; $P = .005$). There was no change in effective hand-off communication between the interns, with experience over time during their internship (using indicator variables representing 2- or 3-month intervals).

In addition, we categorized each important item by type: to-do, anticipa-

tory guidance (if, then), and knowledge. We performed χ^2 tests and found a statistically significant difference in the likelihood that an item would be effectively communicated when it was a to-do item (65%) or an item related to anticipatory guidance (69%) compared with knowledge items (38%; $P = .003$). The number of patients discussed during each hand-off ranged between 3 and 5, with an average of 3.8. We compared the effectiveness of hand-off communication between postcall interns who had fewer than the average number of patients (3 patients) and postcall interns who had more than the average (4 or 5 patients) and found no relationship between the number of hand-off patients and the agreement between the interns on the most important item about a patient (60% vs 55%; $P = .57$).

Postcall interns also overestimated the extent to which on-call interns appreciated the rationale behind the information. When the postcall intern provided a rationale, the on-call intern failed to mention that rationale 60% of the time. In some cases, postcall and on-call interns even provided very different rationales. For example, a postcall intern expected the on-call intern to say that the rationale behind “follow-up with case manager” was to “make sure she talked with patient’s [primary medical doctor],” but the on-call intern actually reported the rationale as “to ensure nothing holding up discharge.”

The average rating of how well prepared the on-call intern felt to take care of patients after the hand-off was 8.8 of 10.0 (SD: 1.0), and the postcall intern reported an average of 8.0 of 10.0 (SD: 1.0). The overall rating of hand-off was 8.3 (SD: 1.3) by the on-call intern and 7.6 (SD: 1.1) by the postcall intern.

DISCUSSION

This study found that pediatric interns overestimate the effectiveness of their hand-off communication, despite their failure to convey the most important information about a patient 40% of the time. This study ties in theories from communication psychology as a possible explanation for why hand-off communication is so poor. In that light, resident miscommunication is the result of a complex interplay among various factors. Because speakers know what they are trying to convey, they tend to think that what they say is clear to anyone.²⁰ Moreover, because they overestimate how well they communicated, postcall interns are less likely to verify whether the on-call intern actually understood^{14,21}; therefore, the inability of the postcall intern to gauge accurately the on-call intern’s understanding of patient information may greatly affect hand-off quality; not only are on-call interns failing to receive important patient information, but also the postcall interns are systematically failing to realize that breakdown of communication. These communication breakdowns occurred even with written hand-offs.

It is interesting that there was no change in effective communication over time. This could mean that increasing clinical knowledge and experience alone do not affect an intern’s ability to communicate effectively during hand-offs. Because there is no formal hand-off curriculum, this suggests that senior residents would not be much better than the interns. It is important to recognize that the literature has found that hand-offs in most residency programs are executed by interns alone with little supervision by senior residents or attendings.⁴ Our findings suggest that even with the presence of a senior resident, postcall interns still overestimate the effectiveness of their hand-off communication.

One possibility for the finding that to-do and anticipatory guidance items were more likely to be communicated compared with knowledge items is that they refer to high-priority items that are relevant for the on-call intern’s upcoming shift. In contrast, items related to knowledge may be less urgent and therefore not as likely to be remembered by the on-call intern. This is in concordance with a review by the National Aeronautics and Space Administration, where hand-off communication has been found to be most effective when it is driven by “problems, hypotheses, and intent” rather than long lists.²² It seems reasonable to train interns to communicate by using this framework and avoid unnecessary knowledge items that are unlikely to be remembered. This will prevent cognitive overload for on-call interns by tailoring information that is communicated. This is an area that requires additional studies.

There are several limitations to this study. It is a single-institution study with a small number of interns, making its generalizability unknown. It is also unclear whether our findings are generalizable to more senior residents and hand-offs in other subspecialties, yet it is reasonable to assume that our findings underestimate the extent of the problem. Hand-offs in our pediatric residency program receive high priority; they occur in a dedicated room and time, have both verbal and written information, and are supervised by senior residents. Despite such good conditions, we found overestimation of hand-off effectiveness by postcall interns. Although there is no survey of hand-off practices in pediatric residency programs, it is known that hand-offs in internal medicine do not always take place in such ideal setups⁴; therefore, a multisite study is likely to find more overestimation and miscommunication.

Unfortunately, some interviews were not conducted because either the postcall or the on-call intern was unavailable. Mainly postcall interns were unable to participate in interviews because they either did not answer a page as a result of workload or left the hospital to adhere to duty hour restrictions. There is no reason to believe that the excluded dyads had more effective hand-offs. If anything, given that the excluded dyads seemed to be under increased pressure, their hand-offs might have been even less effective.

REFERENCES

1. Philibert I, Friedmann P, Williams WT. New requirements for resident duty hours. *JAMA*. 2002;288(9):1112–1114
2. Charap M. Reducing resident work hours: unproven assumptions and unforeseen outcomes. *Ann Intern Med*. 2004;140(10):814–815
3. Steinbrook R. The debate over residents' work hours. *N Engl J Med*. 2002;347(16):1296–1302
4. Horwitz LI, Krumholz HM, Green ML, Huot SJ. Transfers of patient care between house staff on internal medicine wards: a national survey. *Arch Intern Med*. 2006;166(11):1173–1177
5. Landrigan CP, Fahrenkopf AM, Lewin D, et al. Effects of the accreditation council for graduate medical education duty hour limits on sleep, work hours, and safety. *Pediatrics*. 2008;122(2):250–258
6. Institute of Medicine. *Resident Duty Hours: Enhancing Sleep, Supervision, and Safety*. Washington, DC: The National Academies Press; 2008
7. Sutcliffe KM, Lewton E, Rosenthal MM. Communication failures: an insidious contributor to medical mishaps. *Acad Med*. 2004;79(2):186–194
8. Arora V, Johnson J, Lovinger D, Humphrey HJ, Meltzer DO. Communication failures in patient sign-out and suggestions for improvement: a critical incident analysis. *Qual Saf Health Care*. 2005;14(6):401–407
9. White AA, Pichert JW, Bledsoe SH, Irwin C,

CONCLUSIONS

This study shows that postcall interns overestimate the patient information that they convey in hand-offs, and it highlights the extent to which pediatric interns do not agree on the content, priority, or rationale communicated during hand-offs. In the era of restricted duty hours and increased frequency of hand-offs, it is important for educators to consider the role of systematic causes of miscommunication. Future studies should include hand-off improvement efforts, such as the development of specific hand-

off curricula²³ to include emphasis on important items, rationale, and a tailoring of information.²⁴ At the very least, postcall interns should be aware of their “illusion of transparency.” They should appreciate that much of the important information that they thought they conveyed during the hand-off was never really received by the on-call intern.

ACKNOWLEDGMENTS

We thank the residents at the University of Chicago Comer Children's Hospital for willing participation.

- Entman SS. Cause and effect analysis of closed claims in obstetrics and gynecology. *Obstet Gynecol*. 2005;105(5 pt 1):1031–1038
10. White AA, Wright SW, Blanco R, et al. Cause-and-effect analysis of risk management files to assess patient care in the emergency department. *Acad Emerg Med*. 2004;11(10):1035–1041
11. Schoen C, Osborn R, Huynh PT, et al. Taking the pulse of health care systems: experiences of patients with health problems in six countries. *Health Aff (Millwood)*. 2005;(Suppl Web Exclusives):W5–509–W5–525
12. Petersen LA, Brennan TA, O'Neil AC, Cook EF, Lee TH. Does housestaff discontinuity of care increase the risk for preventable adverse events? *Ann Intern Med*. 1994;121(11):866–872
13. Jaggi R, Kitch BT, Weinstein DF, Campbell EG, Hutter M, Weissman JS. Residents report on adverse events and their causes. *Arch Intern Med*. 2005;165(22):2607–2613
14. Keysar B, Henly AS. Speakers' overestimation of their effectiveness. *Psychol Sci*. 2002;13(3):207–212
15. Gilovich T, Savitsky K, Medvec VH. The illusion of transparency: biased assessments of others' ability to read our emotional states. *J Pers Soc Psychol*. 1998;75(2):332–346
16. Vorauer J, Claude S. Perceived versus actual transparency of goals in negotiation. *Pers Soc Psychol Bull*. 1998;24(4):371–385
17. Van Boven L, Gilovich T, Medvec V. The illusion of transparency in negotiations. *Negotiation Journal*. 2003;19(2):117–131
18. Wu S, Keysar B. The effect of information overlap on communication effectiveness. *Cogn Sci*. 2007;31(1):160–181
19. Gillespie R. *Manufacturing Knowledge: A History of the Hawthorne Experiments*. Cambridge, England: Cambridge University Press; 1991
20. Epley N, Keysar B, Van Boven L, Gilovich T. Perspective taking as egocentric anchoring and adjustment. *J Pers Soc Psychol*. 2004;87(3):327–339
21. Keysar B. Communication and miscommunication. *Intercultural Pragmatics*. 2007;4(1):71–84
22. Parke B, Mishkin A. Best practices in shift handover communication: Mars exploration rover surface operations. National Aeronautics and Space Administration. Available at: http://human-factors.arc.nasa.gov/publications/Parke_MER_SurfaceOps_Handovers_05.pdf. Accessed August 15, 2009
23. Horwitz LI, Moin T, Green M. Development and implementation of an oral sign-out skills curriculum. *J Gen Intern Med*. 2007;22(10):1470–1474
24. Patterson ES. Communication strategies from high-reliability organizations: translation is hard work. *Ann Surg*. 2007;245(2):170–172

Interns Overestimate the Effectiveness of Their Hand-off Communication
Vivian Y. Chang, Vineet M. Arora, Shiri Lev-Ari, Michael D'Arcy and Boaz Keysar
Pediatrics 2010;125;491
DOI: 10.1542/peds.2009-0351 originally published online February 8, 2010;

Updated Information & Services

including high resolution figures, can be found at:
<http://pediatrics.aappublications.org/content/125/3/491>

References

This article cites 20 articles, 2 of which you can access for free at:
<http://pediatrics.aappublications.org/content/125/3/491.full#ref-list-1>

Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):
Hospital Medicine
http://classic.pediatrics.aappublications.org/cgi/collection/hospital_medicine_sub
Patient Education/Patient Safety/Public Education
http://classic.pediatrics.aappublications.org/cgi/collection/patient_education:patient_safety:public_education_sub

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<https://shop.aap.org/licensing-permissions/>

Reprints

Information about ordering reprints can be found online:
<http://classic.pediatrics.aappublications.org/content/reprints>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2010 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Interns Overestimate the Effectiveness of Their Hand-off Communication

Vivian Y. Chang, Vineet M. Arora, Shiri Lev-Ari, Michael D'Arcy and Boaz Keysar

Pediatrics 2010;125:491

DOI: 10.1542/peds.2009-0351 originally published online February 8, 2010;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/125/3/491>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since . Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2010 by the American Academy of Pediatrics. All rights reserved. Print ISSN:

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

