

## BC COVID THERAPEUTICS

### Memorandum

Date: April 30, 2021

To: BC Physicians, Nurse Practitioners, Pharmacists and other Healthcare Providers

From: BC COVID-19 Therapeutics Committee & Medical Director, Thrombosis Program, VCH

Re: **Vaccine-induced Immune Thrombotic Thrombocytopenia (VITT)**

**Situation:** Vaccine-induced immune thrombotic thrombocytopenia (VITT), also known as thrombosis with thrombocytopenia syndrome (TTS), is a rare, idiosyncratic life-threatening condition that can arise after COVID-19 vaccination with AstraZeneca (AZ) or Johnson & Johnson/Janssen (JJ) vaccines (not reported with Pfizer or Moderna vaccines). This document with supplemental appendices has been developed through extensive consultation with BC medical experts and physician groups to improve awareness and to help guide screening, diagnosis and management of VITT/TTS.

**Background:** VITT/TTS is a condition characterized by: 1) symptom onset 4 to 28 days after exposure to AZ or JJ vaccine; 2) platelet count less than  $150 \times 10^9/L$ ; and 3) thrombosis, especially in uncommon sites such as cerebral venous sinuses and abdominal veins (eg. portal vein), as well as arterial thrombosis. Mortality is estimated to be approximately 20%.

VITT/TTS resembles heparin-induced thrombocytopenia (HIT) in its mechanism, and diagnosis requires the presence of anti-PF4/heparin antibodies. Similar to HIT, early diagnosis is essential because patients can deteriorate rapidly. Currently, the recommended treatment is empiric initiation of intravenous immunoglobulin (IVIG) and a non-heparin anticoagulant. Based on available data, the best estimate of the risk of VITT/TTS after vaccination with AZ or JJ vaccines is 1 in 100,000 doses. In BC, approximately 400,000 doses of the AZ and JJ vaccines are anticipated to be administered, but this number may increase as our vaccination campaign continues.

**Assessment and Recommendations:** In an effort to best protect British Columbians, BC medical experts have put together 4 documents for dissemination to all physicians and other healthcare providers involved in patient care post-vaccination. These documents (Care Pathway, Primary Care Outpatient Assessment, Key Messages Summary, and a VITT/TTS Poster) aim to educate and support physicians and facilitate rapid diagnosis and treatment of VITT/TTS.

#### *Primary Care Outpatient Assessment*

This provides a concise, point-of-care summary on assessing patients for VITT/TTS in the outpatient office. The emphasis is on vaccine history, high index of suspicion for thrombosis, getting a stat CBC when indicated, and sending a patient to the Emergency department if needed.

#### *Clinical Care Pathway of VITT/TTS in British Columbia*

This algorithm provides guidance to clinicians on how to screen, triage, diagnose and treat patients in a step-wise approach. Not all steps will be relevant to all clinicians, but an understanding of the expected level of care and the required steps is useful. **As the diagnosis of VITT/TTS requires a HIT ELISA test that is only available at St. Paul's Hospital in Vancouver and at the Royal Jubilee Hospital in Victoria, rapid communication with hematologists and laboratory physicians is critical.**

#### *Key Messages and Caveats for VITT/TTS*

These key messages are intended for targeted groups of physicians who will be involved in the care of a patient with suspected or probable VITT/TTS. These statements and recommendations are based on evidence that is rapidly changing, and they should not replace clinician judgement. Case-by-case review with a hematologist or thrombosis specialist is highly recommended.


#### *VITT/TTS Poster*

The poster provides an overview of the main questions regarding VITT/TTS: When should VITT/TTS be suspected? What tests should be ordered? What empiric treatment should be started? This information is useful for all healthcare professionals who may be involved in other aspects of the patients' care.

As new data become available regarding the diagnosis and treatment of VITT/TTS, the BC COVID-19 Therapeutics Committee will provide updated recommendations. Please contact Dr. Agnes Lee, Medical Director, VCH Thrombosis program, at [alee14@bccancer.bc.ca](mailto:alee14@bccancer.bc.ca) for questions.



# Primary Care Outpatient Assessment



**ALL patients presenting with thrombosis symptoms 4-28 days after AZ/JJ vaccine exposure should get a STAT CBC at hospital-based labs or Lifelabs**

## Clinical Care Pathway of VITT/TTS in British Columbia

VITT/TTS Suspected  
AZ or JJ vaccine 4 – 28 days ago AND

- platelet count  $<150 \times 10^9/L$  or
- signs or symptoms of thrombosis

Send to ED for STAT peripheral smear\*, INR, aPTT, D-dimer, fibrinogen and COVID testing  
\*microangiopathy with red cell fragmentation and hemolysis is rarely described in VITT/TTS

Signs of symptoms consistent with **arterial or venous thrombosis ANYWHERE:**

- Severe headache, vision changes or other neurological symptoms
- Unexplained shortness of breath or chest pain
- Unexplained back or abdominal pain
- Swelling or redness in a limb
- Acute pain with pallor in a limb
- Petechiae, easy bruising or bleeding

Any one of:

- High D-dimer
- Low fibrinogen
- Abnormal INR/aPTT
- High suspicion of clot

VITT/TTS Possible

- Ask if recent exposure to heparin/LMWH (within 1-4 weeks)
- Call Hematology (if no local Hematology, call 604-875-5000 to page VGH Hematologist)
- **MUST** call Laboratory to do stat HIT assay for query VITT/TTS (draw 2 red and 2 blue top tubes) and complete requisition<sup>5</sup>
- Order appropriate imaging to look for clots and other labs if indicated

Thrombosis Confirmed or High suspicion of clot

Follow up results for HIT ELISA (Immucor assay at St. Paul's (PHC) or Royal Jubilee (VIHA))

HIT ELISA positive

VITT/TTS Likely  
Start empiric treatment  
Do not wait for lab results

Manage according to standard practice  
Order imaging and labs if needed to rule out thrombosis as dictated by symptoms (eg. CTPA, US, angiogram, ECG, troponin)  
**ONLY CT/MR venogram can rule out CVST**

- Discuss VITT/TTS treatment with Hematology (and Stroke Neurology if CVST confirmed or suspected)
- Give IVIG 0.5 – 1.0 g/kg daily (max total dose 2.0 g/kg) (**Ensure blood drawn for HIT assay BEFORE giving IVIG**)
- Use non-heparin anticoagulant (eg. apixaban, rivaroxaban, fondaparinux, argatroban)
- Consider fibrinogen or cryoprecipitate if patient bleeding and fibrinogen is less than 1.0 g/L
- Do not give platelet unless life-threatening bleeding or need life-saving surgery
- Lab to confirm samples are sent to McMaster for SRA

- Treat as standard VTE/ATE
- DOAC preferred to avoid HIT
- Check platelet count in 3-5 days

VITT/TTS Excluded  
Possible post-vaccine ITP – Contact Hematology

- Legend**
- ATE Arterial thromboembolism
  - AZ AstraZeneca
  - CTPA CT pulmonary angiogram
  - CVST Cerebral venous sinus thrombosis
  - DOAC Direct oral anticoagulant (apixaban, rivaroxaban)
  - ED Emergency Department
  - HIT Heparin-induced thrombocytopenia
  - ITP Immune thrombocytopenic purpura
  - IVIG Intravenous immunoglobulin
  - JJ Johnson & Johnson
  - SRA Serotonin release assay (platelet activation assay)
  - TTS Thrombosis and thrombocytopenia syndrome
  - VITT Vaccine-induced immune thrombotic thrombocytopenia
  - VGH Vancouver General Hospital
  - VTE Venous thromboembolism

# Key Messages and Caveats for VITT/TTS

## Emergency Medicine Primary Care

- Peak period of onset of thrombosis symptoms associated with VITT/TTS is 6 – 14 days after vaccine
- Check if heparin/LMWH exposure within 1 – 4 weeks (if positive, this could be heparin-induced thrombocytopenia [HIT])
- Most cases of VITT present with platelets  $<50 \times 10^9/L$  and D-dimer is typically  $>4x$  ULN
- MUST draw blood work for HIT testing and tell lab to arrange for VITT/TTS testing PRIOR to giving IVIG
- MUST order CT/MRI venogram to rule out cerebral venous sinus thrombosis (CVST) if bad headache or neurological symptoms
- Do not give platelet transfusion unless discussed with Hematology
- Call Hematology if vaccine-associated ITP or evidence of red cell fragments (schistocytes) on smear

## Hematology Internal Med Neurology

- Treat as VITT/TTS unless ruled out with ELISA/SRA (IVIG + use non-heparin anticoagulant +/- steroids)
- If platelets  $<150 \times 10^9/L$  and D-dimer highly elevated, BUT no thrombosis, consider IVIG only or prophylactic anticoagulation while awaiting HIT ELISA results
- If only thrombosis and platelets  $>150 \times 10^9/L$ , start anticoagulant treatment (DOAC preferred) and monitor CBC to exclude early stage of VITT/TTS
- Platelet transfusion only if life-threatening bleeding or life-saving surgery indicated, and only AFTER IVIG has been given
- Consider fibrinogen replacement if less than 1.0 g/L and active bleeding
- Plasma exchange has been used in refractory cases (failure of IVIG and anticoagulant) with variable outcomes

## Laboratory Transfusion Medicine

- Review peripheral smear to rule out other causes of thrombosis + thrombocytopenia
- Sensitivity of HIT ELISA is considered sufficiently high to exclude VITT/TTS (optical density [OD]  $>2.0$  in most cases)
- Non HIT ELISA assays have insufficient sensitivity and unknown specificity
- Serotonin release assay (SRA) at McMaster requires 2 red (or gold) top and 2 blue top tubes and completed lab requisition (<https://fhs.mcmaster.ca/plateletimmunology/vipit.html>)
- Store plasma and serum for all suspected cases (vaccine timing +/- platelets  $<150 \times 10^9/L$  +/- thrombosis)
- Other thrombophilia testing not required nor recommended

# Vaccine-induced Immune Thrombotic Thrombocytopenia (VITT)/ Thrombosis and Thrombocytopenia Syndrome (TTS) Poster

Most important interventions to reduce morbidity and mortality are early diagnosis and empiric IVIG treatment

## When to suspect VITT/TTS?

AZ or JJ vaccine 4 – 28 days prior to onset of thrombosis symptoms

Severe headache, visual changes or other neurological symptoms

Signs or symptoms of other types of clotting (eg. chest pain, leg swelling, abdominal pain)

Unusual bruising, bleeding or petechiae

## What tests to order?

STAT CBC to check platelet count ( $<150 \times 10^9/L$ )

Coagulation tests to check for clotting: D-dimer, INR, aPTT, fibrinogen

HIT ELISA to confirm presence of anti-PF4 antibodies

CT/MRI venogram and other imaging if other sites of clots suspected

## What empiric treatment to start?

DO NOT GIVE PLATELETS

IVIG 0.5 – 1.0 g/kg daily (maximum 2 g/kg total dose)

Non-heparin anticoagulant (eg. direct oral anticoagulant, argatroban)

Contact Hematology or Thrombosis Specialist